



ANTON PANNEKOEK
INSTITUTE

The relativistic jets in the BH XRB MAXI J1836-194

Thomas Russell


J. Miller-Jones, S. Markoff, C. Ceccobello, G. Sivakoff

M. Lucchini, D. Russell, R. Soria, D. Altamirano, A. Tetarenko






UNIVERSITY OF AMSTERDAM


MAXI J1836-194 – a black hole XRB in outburst



JACPOT XRB: Quasi-simultaneous multiwavelength observing campaign






Radio through to X-ray behaviour





AA


Simultaneous broadband data








Broadband evolution




agnjet: an outflow-dominated model




Markoff et al. 2005, 2008
 Gallo et al. 2007
 Maitra et al. 2009, 2011
 Plotkin et al. 2015
 Connors et al. 2017
 Crumley et al. 2017


The inverted radio spectrum




Accounting for the X-ray emission



Fitting the IR and optical




A second jet component?




Fading, flaring?

October 12



October 27



Summary

- Unprecedented multiwavelength coverage
- Evolving compact jet
- Track the jet spectral break
- Broadband modelling hints at two component jet
- Further analysis is ongoing

Conclusions

- Only possible with simultaneous multiwavelength monitoring
- mm and IR – probe inner regions of the jet



Image courtesy of Harvard - SMA



Image courtesy of
ALMA, O. Dessibourg



Image courtesy of ESO

Conclusions

