

The unexpectedly large proportion of high-mass star-forming cores in a Galactic mini-starburst

F. Motte^{1,2*}, T. Nony^{1*}, F. Louvet^{3*}, K. A. Marsh⁴, S. Bontemps⁵, A. P. Whitworth⁴,
A. Men'shchikov², Q. Nguyễn Lương^{6,7}, T. Csengeri⁸, A. J. Maury², A. Gusdorf⁹, E. Chapillon^{5,10},
V. Könyves², P. Schilke¹¹, A. Duarte-Cabral⁴, P. Didelon² and M. Gudel²

Understanding the processes that determine the stellar initial mass function (IMF) is a critical unsolved problem, with profound implications for many areas of astrophysics¹. In molecular clouds, stars are formed in cores—gas condensations sufficiently dense that gravitational collapse converts a large fraction of their mass into a star or small clutch of stars. In nearby star-formation regions, the core mass function (CMF) is strikingly similar to the IMF, suggesting that the shape of the IMF may simply be inherited from the CMF^{2–5}. Here, we present 1.3 mm observations, obtained with the Atacama Large Millimeter/submillimeter Array telescope, of the active star-formation region W43-MM1, which may be more representative of the Galactic-arm regions where most stars form^{6,7}. The unprecedented resolution of these observations reveals a statistically robust CMF at high masses, with a slope that is markedly shallower than the IMF. This seriously challenges our understanding of the origin of the IMF.

The initial mass function (IMF), giving the relative numbers of stars born with different masses, between ~ 0.08 and $\sim 100M_{\odot}$, appears to be universal^{8,9}, although there are a few claims to the contrary, especially in young massive clusters^{10,11}. In nearby star-formation regions, stars with masses between ~ 0.08 and $\sim 5M_{\odot}$ are observed to form in cores, and the core mass function (CMF) in these regions has an approximately log-normal shape with a power-law tail at higher masses ($\gtrsim M_{\odot}$), very similar to the IMF, but with the peak shifted to higher masses by a factor $1/e \sim 2.5 \pm 1$ (refs^{2–5}). This suggests that the shape of the IMF is determined by the processes that determine the CMF, in which case there must be an approximately self-similar mapping from core mass to stellar mass, with a very small variance (see Methods). However, nearby star-formation regions are probably not representative of the regions where most stars are formed and they do not form massive stars (~ 8 to $\sim 100M_{\odot}$)¹². Previous attempts to resolve cores in high-mass star-formation regions have suffered from poor statistics^{13,14} and have therefore not allowed a critical comparison of the CMF and IMF^{14–16}.

W43-MM1 is a massive cloud ($\sim 2 \times 10^4 M_{\odot}$) located at the tip of the Galactic bar—a very rich region in terms of cloud concentration and star-formation activity^{6,7,17} located ~ 5.5 kpc from the Sun¹⁸. Shock tracing lines such as SiO are detected throughout this

cloud, testifying to the prevalence of inflowing and colliding gas streams^{6,19}. The large densities in W43-MM1, traced at high angular resolution by millimetre continuum radiation, suggest it will form a massive star cluster in the near future⁷. We used the Atacama Large Millimeter/submillimeter Array telescope (ALMA) to image the complete W43-MM1 cloud (~ 3 pc²) at 1.3 mm (see Supplementary Fig. 1), with unprecedented resolution ($\sim 2,400$ au), and identified a large sample of individual cores with a broad and previously unmatched range of masses.

Since the 1.3 mm map of W43-MM1 has a dynamic range of 30 in physical scale and 4,000 in flux, we used the *getsources* algorithm²⁰ to identify and characterize the cores. *getsources* decomposes cloud emission into a multi-resolution data-cube; two cube dimensions give the position on the sky, and the third dimension is the physical scale of cloud substructure. Cores were identified at small scales, and their positions, spatial extents (best-fitting ellipses at half-maximum), peak and integrated fluxes (corrected for local background), masses, and mean densities were computed (see Supplementary Table 1). Figure 1 shows the elliptical outlines of the 131 cores returned by *getsources*, of which 94 are deemed to be very robust (significance ≥ 7). The cores have diameters ranging from $\sim 1,300$ to $\sim 3,500$ au once deconvolved with the $0.44''$ (or $\sim 2,400$ au) telescope beam; masses ranging from ~ 1 to $\sim 100M_{\odot}$; and mean H_2 densities ranging from $\sim 10^7$ to $\sim 10^{10}$ cm⁻³. Monte Carlo extraction simulations indicate that the sample is 90% complete down to $\sim 1.6M_{\odot}$ (see Methods).

The blue histogram in Fig. 2a displays the differential CMF of W43-MM1, $dN/d\log(M)$, giving the number of cores in uniformly distributed but variably sized logarithmic bins of mass²¹. The blue histogram in Fig. 2b displays the cumulative CMF of W43-MM1, $N(>\log(M))$, giving the number of cores with M_{core} larger than M . If these histograms are truncated below the 90% completeness limit ($\sim 1.6M_{\odot}$), thereby reducing the sample to 105 cores, they can both be fitted with power laws: $dN/d\log(M) \propto M^{\gamma_{\text{diff}}}$ with $\gamma_{\text{diff}} \simeq -0.90 \pm 0.06$ and $N(>\log(M)) \propto M^{\gamma_{\text{cumul}}}$ with $\gamma_{\text{cumul}} \simeq -0.96 \pm 0.02$ (red lines in Fig. 2a,b). For comparison, the magenta lines in Fig. 2a,b show the slope of the IMF at masses larger than $\sim 1M_{\odot}$, which for both the differential and cumulative forms of the IMF is -1.35 (refs^{9,22}). The range of core masses from ~ 1.6 to $\sim 100M_{\odot}$ corresponds to a range of stellar masses smaller by a factor $1/e \sim 2.5$ (that is, stellar masses

¹Institut de Planétologie et d'Astrophysique de Grenoble, Université Grenoble Alpes, CNRS, Grenoble, France. ²AIM Paris-Saclay/Département d'Astrophysique, CEA, CNRS, Université Paris Diderot, CEA-Saclay, Gif-sur-Yvette, France. ³Department of Astronomy, Universidad de Chile, Santiago, Chile. ⁴School of Physics and Astronomy, Cardiff University, Cardiff, UK. ⁵Laboratoire d'Astrophysique de Bordeaux, OASU, Université Bordeaux, CNRS, Pessac, France. ⁶Korea Astronomy and Space Science Institute, Daejeon, Republic of Korea. ⁷NAOJ Chile Observatory, National Astronomical Observatory of Japan, Tokyo, Japan. ⁸Max-Planck-Institut für Radioastronomie, Bonn, Germany. ⁹LERMA, Observatoire de Paris, PSL Research University, CNRS, Sorbonne Universités, UPMC Université, Paris, France. ¹⁰Institut de Radioastronomie Millimétrique, Saint Martin d'Hères, France. ¹¹Physikalisches Institut, University of Cologne, Cologne, Germany. *e-mail: frederique.motte@univ-grenoble-alpes.fr; thomas.nony@univ-grenoble-alpes.fr; fabien.louvet@gmail.com




Fig. 1 | High-angular-resolution image of the W43-MM1 cloud, revealing a rich population of cores. 1.3 mm dust continuum emission, observed by the ALMA interferometer, is presumed to trace the column density of gas, revealing high-density filaments and embedded cores. The filled yellow ellipse on the right represents the angular resolution, and a scale bar is shown. Ellipses outline core boundaries (at half-maximum) as defined by the getsources²⁰ extraction algorithm. Core masses span the range ~ 1 to $\sim 100 M_{\odot}$ and can therefore be expected to spawn stars with masses from ~ 0.4 to $>40 M_{\odot}$ (ref. 5; see Supplementary Table 1). All cores are shown. Hashed ellipses indicate the most robust identifications.

from ~ 0.6 to $\sim 40 M_{\odot}$, allowing a robust comparison with the higher-mass ($\gtrsim M_{\odot}$) IMF. The result is stable against variations in the temperature model, dust emissivity, extraction algorithm and reduction technique (see Supplementary Table 2). We conclude that at masses larger than $\sim 1.6 M_{\odot}$, the CMF in W43-MM1 is markedly flatter than the IMF. This result seriously challenges the widespread assumption that the shape of the IMF is inherited directly from the CMF.

The main uncertainty in the CMF derives from the estimation of core masses using their measured 1.3 mm continuum fluxes (see Methods). For the most massive cores, we used a 1.3 mm image based on the signal in a narrow composite band (~ 65 MHz wide) that is not contaminated by line emission. For the lower-mass cores (the majority), we used an image based on the signal in the full band (~ 1.9 GHz wide), since line contamination for low-mass cores is expected to be negligible (see C_{line} in Supplementary Table 1). Variations in dust emissivity are presumed to be small among the W43-MM1 cores, since 90% of them have uniformly high mean densities ($\bar{n}_{\text{H}_2} \sim 3 \times 10^7 - 3 \times 10^8 \text{ cm}^{-3}$) and warm temperatures ($\bar{T}_{\text{dust}} \sim 23 \pm 2 \text{ K}$) (see Supplementary Table 1). The main source of uncertainty is the dust temperature, since this is critical for converting flux into mass. Figure 3 displays the mean dust temperatures used for estimating core masses. A mean line-of-sight dust temperature was estimated using Herschel 70–500 μm images⁶, Atacama Pathfinder Experiment 350 and 870 μm images¹⁸, and mosaics obtained with ALMA (present data) and Institute for Radio Astronomy in the Millimeter range (IRAM)⁷ interferometers at 1.3 and 3 mm, respectively. By applying the Bayesian point process mapping (PPMAP) procedure²³, we obtained column-density maps in different dust-temperature slices. The mean dust temperature in each pixel is then a column-density-weighted average along that line of sight. However, in the vicinity of 10 hot cores (of which 3 have been identified previously^{24,25}), the local heating is not properly traced by the 2.5''-resolution PPMAP temperature image. Here, we divide the total luminosity of the W43-MM1 cloud ($\sim 2 \times 10^4 L_{\odot}$) between the cores in proportion to their associated line contamination in the wide 1.3 mm band. We then use the individual luminosities, $L_{\star} < 10$ to $\sim 10^4 L_{\odot}$, and PPMAP background temperatures to esti-




Fig. 2 | W43-MM1 CMFs challenging the relationship between the CMF and IMF. **a, b**, Differential (**a**) and cumulative form (**b**). Above the sample 90% completeness limit, estimated to be $M_{\text{core}} = 1.6 M_{\odot}$ (black vertical line), the W43-MM1 CMFs (blue histograms) are well fit by the single power laws $dN/d\log(M) \propto M^{-0.90}$ in **a** and $N(>\log(M)) \propto M^{-0.96}$ in **b** (red lines and 1σ uncertainties). The error bars on the differential CMF correspond to \sqrt{N} counting statistics. The cumulative CMF in **b** is the more robust statistically: its 5σ global uncertainty (± 0.13 , hatched area) was estimated from Monte Carlo simulations. The W43-MM1 CMF is clearly flatter than the IMF^{9,22}, which in the corresponding mass range has slopes $dN/d\log(M) \propto M^{-1.35}$ and $N(>\log(M)) \propto M^{-1.35}$ (magenta lines).

mate the mean core dust temperatures ($\bar{T}_{\text{dust}} \sim 20$ to $\sim 90 \text{ K}$) and their uncertainties (see Supplementary Table 1) using an approximate radiation transport model²⁶. Monte Carlo simulations indicate that the mass uncertainties in Supplementary Table 1 correspond to a 5σ uncertainty of ± 0.13 in the slope of the CMF (see Methods and black-hatched area in Fig. 2b).

The fidelity of the CMF also depends on whether we have correctly identified cores; that is, structures that (1) are gravitationally bound and are therefore destined to spawn stars and (2) already contain most of the mass that will eventually end up in those stars. Parenthetically, we note that core masses probably grow with time due to inflowing gas streams such as those observed towards many massive cores^{27–29}. In respect of criterion (1), studies of gravitational boundedness using ¹³CS (5–4) lines from the present ALMA project to determine internal velocity dispersions indicate that W43-MM1 cores with $M_{\text{core}} > 12 M_{\odot}$ are secure but the status of lower-mass cores would benefit from further investigation. In respect of criterion (2), the low luminosity-to-mass ratio of the whole region, $L_{\text{bol}}/M \sim 5 L_{\odot}/M_{\odot}$ (ref. 17), and the low mean temperature, $\bar{T}_{\text{dust}} \sim 20 \text{ K}$ (see Fig. 3), imply that the region is young. We note that the two most massive cores may assimilate further mass from their dense surroundings and eventually form stars of $\sim 100 M_{\odot}$, with $L_{\star} \sim 10^5 L_{\odot}$ on the main sequence. However, currently they are 10–50 times less luminous. We conclude that any protostars embedded within the W43-MM1 cores are at the very beginning of their accretion phase and therefore contain only a small fraction of their final stellar mass.

Finally, the fidelity of the CMF depends on the completeness of the core sample. Owing to increased source and background




Fig. 3 | Mean estimated dust temperatures for cores. Over most of the frame, \bar{T}_{dust} is the column-density-weighted mean obtained with PPMAP²³ using $2.5''$ (or 14,000 au) resolution (colour scale). In the vicinity of the 10 most luminous sources, the mean core temperature with $0.44''$ (or 2,400 au) resolution was calculated using an approximate radiation transport model based on their estimated luminosities²⁶. The W43-MM1 filaments are located by the 4, 20 and 50 mJy beam⁻¹ contours of the 1.3 mm continuum map of Fig. 1. The \bar{T}_{dust} map shows cloud heating by the nearby cluster, screening by the W43-MM1 high-density filaments and strong local heating by the 10 most luminous (30 to $10^4 L_{\odot}$) cores.

confusion in the denser parts of the cloud, the core sample is 90% complete above $\sim 1.6 M_{\odot}$ outside the main filament and above $\sim 4.5 M_{\odot}$ within it (see Supplementary Fig. 2). As in previous CMF evaluations, we use the median of the detection thresholds (here $\sim 1.6 M_{\odot}$), but even with the more conservative threshold of $\sim 4.5 M_{\odot}$, the CMF is still markedly shallower than the IMF (see Supplementary Table 2). The 5σ uncertainty in the fit to the CMF on Fig. 2b (black-hatched area) reflects the above assumptions, along with contributions from the data-reduction method, extraction algorithm²⁹, and effects of source and background confusion (see Methods).

Owing to the high sensitivity and resolution of our ALMA 1.3 mm image of the W43-MM1 cloud, we were able to obtain a robust CMF covering masses in the range ~ 1.6 to $\sim 100 M_{\odot}$, and hence to conclude that the CMF in W43-MM1 is much shallower than the higher-mass ($\gtrsim M_{\odot}$) IMF; namely, $N(>\log(M)) \propto M^{\gamma}$ with $\gamma \simeq -0.96 \pm 0.13$ instead of $\gamma \simeq -1.35$. This is in stark contrast with previous robust evaluations of the CMF in other regions and at lower masses, where the CMF appears to be very similar to the IMF²⁻⁵. Scenarios that might explain our result and still allow the shape of the IMF to be inherited from the shape of the CMF, with an approximately self-similar mapping, fall into two broad categories. In the first, W43-MM1 is not representative of the environments in which most low-mass stars form^{10,11}; to obtain the complete IMF, stars formed in environments like W43-MM1 must be mixed with stars formed in environments that contain a higher proportion of low-mass cores. However, this scenario is hard to justify if the local IMF observed in extreme star-formation regions is similar to the universal IMF⁸. In the second, massive cores are over-represented in W43-MM1; either massive cores live significantly longer than lower-mass cores³⁰ (but current lifetime derivations¹² suggest the opposite) or star formation in regions like W43-MM1 is prolonged, and massive cores and the stars they spawn only form during a short period (probably near the beginning), while lower-mass cores and low-mass stars form over a much longer period (which seems unlikely because it would require the low-mass period to be nearly ten times longer than the high-mass period, although this cannot

be ruled out). If none of the above scenarios appears to explain our results, that would imply that the mapping from the CMF to the IMF is not statistically self-similar. Either higher-mass cores must convert a smaller fraction of their mass into stars than lower-mass cores (which seems unlikely¹⁴) or they must spawn more stars with a wider logarithmic range of masses (which seems more likely), or some combination of these possibilities exists. The shape of the IMF—at least at masses greater than $\sim M_{\odot}$ —is then not simply inherited from the shape of the CMF and the processes that determine the IMF remain to be determined.

Methods

Observations and data reduction. The W43-MM1 ALMA observations were carried out between September 2014 and June 2015 (project number 2013.1.01365.S) using the 12 m interferometric array. With baselines ranging from 13 to 1,045 m, the 1.3 mm image is sensitive to emission on angular scales from the synthesized beam ($0.37'' \times 0.53''$ ($\sim 2,400$ AU at 5.5 kpc)) to the filtering scale of the interferometer for our observations ($12''$). The W43-MM1 cloud was imaged with a 33-field mosaic covering its $2.1 \text{ pc} \times 1.4 \text{ pc}$ extent. The complete dataset contains eight spectral windows, of which the one most used here is dedicated to the 1.3 mm continuum centred at 233.45 GHz, with a bandwidth of 1.9 GHz and a channel step of 977 kHz. The complementary data taken with the 7 m interferometric array were not used because of their low signal-to-noise ratio.

Data were reduced using the CASA 4.3.1 software³¹, applying manual and SelfCalibration scripts. The SelfCalibration technique improved image quality provided that it started from the highest-sensitivity map integrated over the complete 1.9 GHz continuum band. We then applied the Clean algorithm with a robust weight of 0.5 and the MultiScale option³², which minimizes interferometric artefacts associated with missing short spacings. We did not use merged 12 and 7 m data because the MultiScale cleaning of 12 m-only data provides the optimum image sensitivity for extracting cores. From the residual map of the continuum map integrated over the complete 1.9 GHz band, we estimated noise levels from $\sim 0.13 \text{ mJy beam}^{-1}$ in the outskirts to $\sim 2.5 \text{ mJy beam}^{-1}$ towards the main filament. This variation was due to confusion caused by strong, extended and structured cloud emission³³.

Core extraction. Compact sources were extracted using getsources (version 1.140127)—a multi-scale, multi-wavelength source-extraction algorithm developed for Herschel CMF studies^{30,34}. The main advantage of this algorithm applied to interferometric images is that it can extract sources in varying backgrounds, from strong filaments to negative areas associated with missing short spacings. We also used the multi-wavelength design of getsources to simultaneously treat the highest-sensitivity (1.9 GHz-integrated) and line-free (65 MHz-integrated) 1.3 mm continuum images. During the detection step, getsources combines both images and defines a catalogue of sources with unique positions. At the measurement stage, getsources measures the background and flux of these sources on each map independently.

We post-processed the getsources catalogue to remove, through visual inspection, sources that were too extended, whose ellipticity was too large to correspond to cores or that were not centrally peaked. The final catalogue contains 131 sources with significance >5 (reliable), of which 94 cores have significance >7 (robust). Core characteristics were measured in the highest-sensitivity continuum image, except when core fluxes in the highest-sensitivity map were larger than those measured in the line-free image, due to line contamination. Sources with line contamination were massive cores harbouring hot cores. Supplementary Table 1 lists the cores detected in W43-MM1, and provides their significance, coordinates, diameters, line contamination ratio, and peak and total integrated flux (corrected for line contamination and local background). While the significance variable reflects the detection significance (equivalent to the signal-to-noise ratio of a source on the scale where it is best detected), flux uncertainties reflect the quality of the flux measurements.

Core mass estimates. The total mass of a core (gas plus dust), having uniform opacity throughout its solid angle, is given by

$$\begin{aligned}
 M_{\text{core}} &= -\frac{\Omega_{\text{beam}} d^2}{\kappa_{1.3\text{mm}}} \ln \left(1 - \frac{S_{1.3\text{mm}}^{\text{peak}}}{\Omega_{\text{beam}} B_{1.3\text{mm}}(T_{\text{dust}})} \right) \times \frac{S_{1.3\text{mm}}^{\text{int}}}{S_{1.3\text{mm}}^{\text{peak}}} \\
 &= -M_{\text{core}}^{\text{opt thin}} \times \frac{\Omega_{\text{beam}} B_{1.3\text{mm}}(T_{\text{dust}})}{S_{1.3\text{mm}}^{\text{peak}}} \\
 &\quad \ln \left(1 - \frac{S_{1.3\text{mm}}^{\text{peak}}}{\Omega_{\text{beam}} B_{1.3\text{mm}}(T_{\text{dust}})} \right)
 \end{aligned}$$

Here, Ω_{beam} is the solid angle of the beam, $d = 5.5 \text{ kpc}$ is the distance to the core, $\kappa_{1.3\text{mm}}$ is the dust opacity per unit mass of gas and dust at 1.3 mm, $S_{1.3\text{mm}}^{\text{peak}}$ and

$S_{1.3\text{mm}}^{\text{int}}$ are the peak and integrated monochromatic fluxes of the core at 1.3 mm, and $B_{1.3\text{mm}}(T_{\text{dust}})$ is the Planck function at dust temperature T_{dust} . $M_{\text{core}}^{\text{opt thin}} = S_{1.3\text{mm}}^{\text{int}} d^2 / (\kappa_{1.3\text{mm}} B_{1.3\text{mm}}(T_{\text{dust}}))$ is the core mass in the optically thin limit. We have adopted $\kappa_{1.3\text{mm}} = 0.01 \text{ cm}^2 \text{ g}^{-1}$ (ref. 35) as being most appropriate for high-density, cool to warm cores. Supplementary Table 1 lists for each core its mass and mean number density³⁶.

The 1.3 mm continuum flux of a core arises mostly from thermal dust emission, which is optically thin except towards the seven densest, most massive cores. Contamination by free-free emission is estimated to be low (<20%; ref. 17) for W43-MM1 cores, since none of the embedded protostars are sufficiently hot and luminous (<10³L_⊙ for all except 3 cores) to ionize significant H II regions. Line contamination was evaluated by comparing the fluxes measured in the total 1.9 GHz continuum band and in a selection of line-free channels summing up to 65 MHz. The mass uncertainties were derived from the flux uncertainties measured by getsources, and from the estimated dust-temperature error. We added the ±1 K errors of PPMAP temperatures²³ to the uncertainty derived from the self-shielding of 0.5'' starless cores and/or the internal heating of 0.5'' protostellar cores lying within the 2.5'' PPMAP resolution^{26,37}. We estimated the absolute values of the core masses to be uncertain by a factor of a few, and the relative values between cores to be uncertain by ~50%.

To estimate the sample completeness, we performed 10 Monte Carlo simulations that placed 2,000 synthetic cores with masses in the range 1–10M_⊙ on the worst-sensitivity background image determined by getsources or on the original image containing real cores. Like the observed cores, the synthetic cores had small sizes, and this reduced confusion by nearby sources and gave an excellent measurement of the total core mass. We then simulated observation of these synthetic protoclusters and applied the same core extraction process as for the real data. Supplementary Fig. 2 shows that the 90% completeness level depends weakly on the background intensity. It is $\sim 0.8 \pm 0.1 M_{\odot}$ in the outskirts of the protocluster, $\sim 1.6 \pm 0.1 M_{\odot}$ where most cores lie and up to $\sim 4.5 \pm 1 M_{\odot}$ on the main filament (see the areas outlined in Supplementary Fig. 1). The 90% completeness level also depends weakly on the source confusion, with an increase up to $2.4 \pm 0.2 M_{\odot}$ for overlapping sources, relative to the mean value, which remains at $\sim 1.6 M_{\odot}$ (see Supplementary Fig. 2).

CMFs. We performed numerous tests and Monte Carlo simulations to prove the robustness of the CMF shape against interferometric artefacts, extraction methods (getsources²⁰ or MRE-GaussClumps²⁹), mass estimates and CMF representations. In developing the best strategy to reduce the W43-MM1 ALMA image, the core catalogue steadily improved, with progressively fewer false cores and more solar-type cores. The 94 robust cores (>7σ) were detected in almost all runs. Supplementary Table 2 lists the major tests performed to evaluate the uncertainty of the CMF fit. For statistical reasons, fits were more robust in cumulative form³⁸, with the complete sample of 131 reliable cores and the complete 1.6–100M_⊙ mass range. However, Supplementary Table 2 shows that the higher-mass part (>4.5M_⊙) of the W43-MM1 CMF is still always flatter than the IMF. Monte Carlo simulations (with 100,000 runs) showed that the mass uncertainties corresponded to the slope of the CMF, γ_{cumul} , having a 5σ uncertainty of ±0.13. Building separate mass functions for cores that do and do not contain accreting protostars was not feasible here. First, the most massive cores all contained young low-luminosity accreting protostars. Second, their outflows made it very hard to ascertain the precise nature of their lower-mass neighbours (that is, whether or not they too already contained accreting protostars).

Self-similar mapping. A statistically self-similar mapping means that, for example, the probability that a core with mass in the range 1.0–1.1M_⊙ spawns—possibly along with other stars—a star with mass in the range 0.40–0.44M_⊙ is the same as the probability that a core with mass in the range 10–11M_⊙ spawns a star with mass in the range 4.0–4.4M_⊙. A mapping with small variance is required because, if the variance were large, the peak of the IMF would be noticeably broader than the peak of the CMF, and this does not appear to be the case³. (We note that incompleteness at low core masses can only make the peak of the CMF broader than current estimates and thereby strengthen this constraint).

To compute a notional conversion efficiency (the mean of the total mass of stars subsequently spawned by a core divided by the core's mass), we need to determine: (1) the mean factor by which a core's mass grows between the time when it is measured to determine the CMF and the time when it has finished forming stars, $\mu_{\text{map}} (>1)$; and (2) the mean fraction of this final mass that goes into stars, $\eta_{\text{map}} (<1)$. The notional efficiency is then $\mu_{\text{map}}\eta_{\text{map}}$ and can in principle exceed unity (which is why it is 'notional'). If we also know (3) the mean number of stars spawned by a single core, N_{map} , we can compute the upwards shift from the peak of the IMF to the peak of the CMF, $1/\epsilon = N_{\text{map}} / (\mu_{\text{map}}\eta_{\text{map}})$. Finally, if we know (4) the logarithmic variance of the mean distribution of stellar masses spawned by a single core, σ_{map} , this must be convolved with the logarithmic variance of the CMF to determine the logarithmic variance of the IMF, $\sigma_{\text{IMF}}^2 = \sigma_{\text{CMF}}^2 + \sigma_{\text{map}}^2$. Since σ_{map} is necessarily finite, a self-similar mapping requires $\sigma_{\text{IMF}} > \sigma_{\text{CMF}}$.

Data availability. The data that support the plots within this paper and other findings of this study are available from the corresponding authors (E.M., T.N. or F.L.) upon reasonable request. Requests for materials should be addressed to E.M., T.N. or F.L.

Received: 3 May 2017; Accepted: 26 March 2018;
Published online: 30 April 2018

References

- Offner, S. S. R. et al. in *Protostars and Planets VI* (eds Beuther, H., Klessen, R. S., Dullemond, C. P. & Henning, Th.) 53–75 (Univ. Arizona Press, Tucson, 2014).
- Motte, F., André, P. & Neri, R. The initial conditions of star formation in the Rho Ophiuchi main cloud: wide-field millimeter continuum mapping. *Astron. Astrophys.* **336**, 150–172 (1998).
- Testi, L. & Sargent, A. I. Star formation in clusters: a survey of compact millimeter-wave sources in the Serpens Core. *Astrophys. J. Lett.* **508**, L91–L94 (1998).
- Enoch, M. L. et al. The mass distribution and lifetime of prestellar cores in Perseus, Serpens, and Ophiuchus. *Astrophys. J.* **684**, 1240–1259 (2008).
- Könyves, V. et al. A census of dense cores in the Aquila cloud complex: SPIRE/PACS observations from the Herschel Gould Belt survey. *Astron. Astrophys.* **584**, A91 (2015).
- Nguyen Luong, Q. et al. Low-velocity shocks traced by extended SiO emission along the W43 ridges: witnessing the formation of young massive clusters. *Astrophys. J.* **775**, 88 (2013).
- Louvet, F. et al. The W43-MM1 mini-starburst ridge, a test for star formation efficiency models. *Astron. Astrophys.* **570**, A15 (2014).
- Bastian, N., Covey, K. R. & Meyer, M. R. A universal stellar initial mass function? A critical look at variations. *Annu. Rev. Astron. Astrophys.* **48**, 339–389 (2010).
- Kroupa, P. et al. in *Planets, Stars and Stellar Systems* Vol. 5 (eds Oswalt, T. D. & Gilmore, G.) 115 (Springer, Dordrecht, 2013).
- Harayama, Y., Eisenhauer, F. & Martins, F. The initial mass function of the massive star-forming region NGC 3603 from near-infrared adaptive optics observations. *Astrophys. J.* **675**, 1319–1342 (2008).
- Maia, F. F. S., Moraux, E. & Joncour, I. Young and embedded clusters in Cygnus-X: evidence for building up the initial mass function? *Mon. Not. R. Astron. Soc.* **458**, 3027–3046 (2016).
- Motte, F., Bontemps, S. & Louvet, F. High-mass star and massive cluster formation in the Milky Way. *Annu. Rev. Astron. Astrophys.* <https://doi.org/10.1146/annurev-astro-091916-055235> (2018).
- André, P. et al. From filamentary clouds to prestellar cores to the stellar IMF: initial highlights from the Herschel Gould Belt survey. *Astron. Astrophys.* **518**, L102 (2010).
- Bontemps, S., Motte, F., Csengeri, T. & Schneider, N. Fragmentation and mass segregation in the massive dense cores of Cygnus X. *Astron. Astrophys.* **524**, A18 (2010).
- Zhang, Q., Wang, K., Lu, X. & Jiménez-Serra, I. Fragmentation of molecular clumps and formation of a protocluster. *Astrophys. J.* **804**, 141 (2015).
- Cheng, Y. et al. The core mass function in the massive protocluster G286.21+0.17 revealed by ALMA. *Astrophys. J.* **853**, 160 (2018).
- Motte, F., Schilke, P. & Lis, D. C. From massive protostars to a giant H II region: submillimeter imaging of the Galactic ministarburst W43. *Astrophys. J.* **582**, 277–291 (2003).
- Nguyen Luong, Q. et al. W43: the closest molecular complex of the Galactic bar? *Astron. Astrophys.* **529**, A41 (2011).
- Louvet, F. et al. Tracing extended low-velocity shocks through SiO emission. Case study of the W43-MM1 ridge. *Astron. Astrophys.* **595**, A122 (2016).
- Men'shchikov, A. et al. A multi-scale, multi-wavelength source extraction method: getsources. *Astron. Astrophys.* **542**, A81 (2012).
- Maíz Apellániz, J. & Úbeda, L. Numerical biases on initial mass function determinations created by binning. *Astrophys. J.* **629**, 873–880 (2005).
- Salpeter, E. E. The luminosity function and stellar evolution. *Astrophys. J.* **121**, 161 (1955).
- Marsh, K. A., Whitworth, A. P. & Lomax, O. Temperature as a third dimension in column-density mapping of dusty astrophysical structures associated with star formation. *Mon. Not. R. Astron. Soc.* **454**, 4282–4292 (2015).
- Herpin, F. et al. The massive protostar W43-MM1 as seen by Herschel-HIFI water spectra: high turbulence and accretion luminosity. *Astron. Astrophys.* **542**, A76 (2012).
- Sridharan, T. K. et al. Hot core, outflows, and magnetic fields in W43-MM1 (G30.79 FIR 10). *Astrophys. J. Lett.* **783**, L31 (2014).
- Motte, F. & André, P. The circumstellar environment of low-mass protostars: a millimeter continuum mapping survey. *Astron. Astrophys.* **365**, 440–464 (2001).
- Hatchell, J. & Fuller, G. A. Star formation in Perseus. IV. Mass-dependent evolution of dense cores. *Astron. Astrophys.* **482**, 855–863 (2008).

28. Swift, J. J. & Williams, J. P. On the evolution of the dense core mass function. *Astrophys. J.* **679**, 552–556 (2008).
29. Csengeri, T. et al. The ATLASGAL survey: a catalog of dust condensations in the Galactic plane. *Astron. Astrophys.* **565**, A75 (2014).
30. Clark, P. C., Klessen, R. S. & Bonnell, I. A. Clump lifetimes and the initial mass function. *Mon. Not. R. Astron. Soc.* **379**, 57–62 (2007).
31. McMullin, J. P., Waters, B., Schiebel, D., Young, W. & Golap, K. CASA architecture and applications. In *Astronomical Data Analysis Software and Systems XVI* (eds Shaw, R. A., Hill, F. & Bell, D. J.) **376**, 127 (Astronomical Society of the Pacific, 2007).
32. Rau, U. & Cornwell, T. J. A multi-scale multi-frequency deconvolution algorithm for synthesis imaging in radio interferometry. *Astron. Astrophys.* **532**, A71 (2011).
33. Hennemann, M. et al. The spine of the swan: a Herschel study of the DR21 ridge and filaments in Cygnus X. *Astron. Astrophys.* **543**, L3 (2012).
34. Men'shchikov, A. A multi-scale filament extraction method: getfilaments. *Astron. Astrophys.* **560**, A63 (2013).
35. Ossenkopf, V. & Henning, T. Dust opacities for protostellar cores. *Astron. Astrophys.* **291**, 943–959 (1994).
36. Tigé, J. et al. The earliest phases of high-mass star formation, as seen in NGC 6334 by Herschel-HOBYS. *Astron. Astrophys.* **602**, A77 (2017).
37. Bracco, A. et al. Probing changes of dust properties along a chain of solar-type prestellar and protostellar cores in Taurus with NIKA. *Astron. Astrophys.* **604**, A52 (2017).
38. Reid, M. A. & Wilson, C. D. High-mass star formation. III. The functional form of the submillimeter clump mass function. *Astrophys. J.* **650**, 970–984 (2006).

Acknowledgements

This paper makes use of the following ALMA data: #2013.1.01365.S. ALMA is a partnership of ESO (representing its member states), NSF (USA) and NINS (Japan), together with NRC (Canada), MOST and ASIAA (Taiwan), and KASI (Republic of Korea), in cooperation with the Republic of Chile. The Joint ALMA Observatory is

operated by ESO, AUI/NRAO and NAOJ. This project has received funding from the European Union's Horizon 2020 research and innovation programme StarFormMapper under grant agreement number 687528. This work was supported by the Programme National de Physique Stellaire and Physique et Chimie du Milieu Interstellaire of CNRS/INSU (with INC/INP/IN2P3), co-funded by CEA and CNES. A.P.W. gratefully acknowledges the support of a consolidated grant (ST/K00926/1) from the UK Science and Technology Funding Council. T.C. acknowledges support from the Deutsche Forschungsgemeinschaft via the SPP (priority programme) 1573 'Physics of the ISM'. A.J.M. has received funding from the European Research Council under the European Union's Horizon 2020 research and innovation programme (MagneticYSOs, grant agreement number 679937).

Author contributions

F.M. and F.L. led the project. E.C., T.N., F.M. and A.J.M. reduced the ALMA data. F.L. ran getsources and the CASA simulator. T.C. ran MRE-GaussClumps. K.A.M. ran PPMAP. S.B. and A.M. performed the Monte Carlo simulations. F.M., T.N. and F.L. analysed the CMF results. F.M. and A.P.W. wrote the manuscript. F.M., S.B., F.L., Q.N.L., A.J.M. and P.S. contributed to the ALMA proposal. All authors discussed the results and implications and commented on the manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary information is available for this paper at <https://doi.org/10.1038/s41550-018-0452-x>.

Reprints and permissions information is available at www.nature.com/reprints.

Correspondence and requests for materials should be addressed to F.M. or T.N. or F.L.


Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

In the format provided by the authors and unedited.


The unexpectedly large proportion of high-mass star-forming cores in a Galactic mini-starburst

F. Motte^{1,2*}, T. Nony^{1*}, F. Louvet^{3*}, K. A. Marsh⁴, S. Bontemps⁵, A. P. Whitworth⁴,
A. Men'shchikov², Q. Nguyễn Lương^{6,7}, T. Csengeri⁸, A. J. Maury², A. Gusdorf⁹, E. Chapillon^{5,10},
V. Könyves², P. Schilke¹¹, A. Duarte-Cabral⁴, P. Didelon² and M. Gaudel²

¹Institut de Planétologie et d'Astrophysique de Grenoble, Université Grenoble Alpes, CNRS, Grenoble, France. ²AIM Paris-Saclay/Département d'Astrophysique, CEA, CNRS, Université Paris Diderot, CEA-Saclay, Gif-sur-Yvette, France. ³Department of Astronomy, Universidad de Chile, Santiago, Chile. ⁴School of Physics and Astronomy, Cardiff University, Cardiff, UK. ⁵Laboratoire d'Astrophysique de Bordeaux, OASU, Université Bordeaux, CNRS, Pessac, France. ⁶Korea Astronomy and Space Science Institute, Daejeon, Republic of Korea. ⁷NAOJ Chile Observatory, National Astronomical Observatory of Japan, Tokyo, Japan. ⁸Max-Planck-Institut für Radioastronomie, Bonn, Germany. ⁹LERMA, Observatoire de Paris, PSL Research University, CNRS, Sorbonne Universités, UPMC Université, Paris, France. ¹⁰Institut de Radioastronomie Millimétrique, Saint Martin d'Hères, France. ¹¹Physikalisches Institut, University of Cologne, Cologne, Germany. *e-mail: frederique.motte@univ-grenoble-alpes.fr; thomas.nony@univ-grenoble-alpes.fr; fabien.louvet@gmail.com



Supplementary Figure 1 — High-angular resolution image of the complete W43-MM1 cloud. 1.3 mm dust continuum emission, observed by the ALMA interferometer, is presumed to trace the column density of gas, revealing high-density filaments and embedded cores. The filled yellow ellipse on the left shows the angular resolution, and a scale bar is shown on the right. Contours correspond to the $S_{1.3\text{mm}}^{\text{peak}} = 4$ mJy/beam flux level (black) and the 10^{23} cm⁻² column density level from *Herschel* (white). Despite the high-sensitivity of the ALMA image, and the high column densities observed, there appear to be no filaments or cores, with a completeness level of $\sim 0.8 M_{\odot}$, in the outskirts of the cloud. The core sample is 90% complete down to $\sim 1.6 M_{\odot}$ on average, outside the main filament (between the two contours); and complete down to $\sim 4.5 M_{\odot}$ within it.



Supplementary Figure 2 — Completeness level of the core sample of Supplementary Table 1. We have performed Monte-Carlo simulations to investigate the sample completeness level as a function of background intensity (red and black curves) and source confusion (blue curve). Synthesized cores are placed (a) on the worst-sensitivity background image, which was determined by *getsources* after subtracting cores of Supplementary Table 1 from Fig. 1, for background confusion tests; and (b) on the original image containing real cores, for source confusion tests. The observations are then simulated and the cores extracted as for the real data. Confusion, due to either a strong background, or the proximity of bright sources, is minimized by the multi-scale approach of *getsources*. Simulations show that the core sample of Supplementary Table 1 is 90% complete down to $\sim 1.6 \pm 0.1 M_{\odot}$ (red curve) and to $\sim 4.5 \pm 1 M_{\odot}$ (black curve) in, respectively, the median- and high-background areas outlined in Supplementary Fig. 1. The census of cores overlapping with others remains 90% complete down to $\sim 2.4 \pm 0.2 M_{\odot}$ (blue curve).

Supplementary Table 1 — Characteristics of the cores detected at 1.3 mm in W43-MM1.

#	RA [J2000]	Dec. [J2000]	Size ["×"]	FWHM [AU]	Sig	C_{line} [%]	$S_{1.3\text{mm}}^{\text{peak}}$ [mJy/beam]	$S_{1.3\text{mm}}^{\text{int}}$ [mJy]	\bar{T}_{dust} [K]	M_{core} [M_{\odot}]	\bar{n}_{H_2} $\times 10^7 \text{cm}^{-3}$
1	18:47:47.02	-1:54:26.86	0.82 × 0.45	2300	1100	45	277 ± 5	592 ± 6	74 ± 2	102 ± 5	210 ± 10
2	18:47:46.84	-1:54:29.30	0.48 × 0.45	1200	670	26	192 ± 5	261 ± 6	59 ± 4	55 ± 6	750 ± 80
3	18:47:46.37	-1:54:33.41	0.52 × 0.47	1200	330	13	109 ± 2	222 ± 2	45 ± 1	59 ± 2	760 ± 30
4	18:47:46.98	-1:54:26.42	0.88 × 0.45	2500	340	74	109 ± 5	312 ± 5	88 ± 7	36 ± 3	57 ± 5
5	18:47:46.76	-1:54:31.21	0.55 × 0.45	1300	170	15	52 ± 4	78 ± 5	47 ± 1	18 ± 1	210 ± 20
6	18:47:46.16	-1:54:33.30	0.55 × 0.45	1300	130	0	46.8 ± 0.8	94 ± 1	23 ± 2	56 ± 9	660 ± 100
7	18:47:47.26	-1:54:29.70	0.48 × 0.45	1200	130	0	47 ± 1	59 ± 1	30 ± 2	23 ± 2	320 ± 30
8	18:47:46.54	-1:54:23.15	0.54 × 0.45	1200	130	0	45 ± 2	61 ± 2	45 ± 5	14 ± 2	190 ± 30
9	18:47:46.48	-1:54:32.54	0.63 × 0.45	1600	110	17	37 ± 2	87 ± 3	50 ± 1	17.8 ± 0.9	97 ± 5
10	18:47:46.91	-1:54:29.99	0.67 × 0.5	2100	91	24	35 ± 4	78 ± 5	51 ± 2	16 ± 1	43 ± 4
11	18:47:46.52	-1:54:24.26	0.52 × 0.45	1200	72	60	16 ± 2	21 ± 2	93 ± 11	2.1 ± 0.3	28 ± 5
12	18:47:46.57	-1:54:32.04	0.72 × 0.45	2000	77	0	30 ± 2	57 ± 2	23 ± 2	31 ± 4	100 ± 10
13	18:47:46.92	-1:54:28.62	0.69 × 0.45	1900	63	0	22 ± 6	22 ± 6	24 ± 2	11 ± 3	40 ± 10
14	18:47:46.97	-1:54:29.66	0.72 × 0.45	2000	57	0	23 ± 6	33 ± 6	22 ± 2	19 ± 4	60 ± 10
15	18:47:44.77	-1:54:45.22	0.66 × 0.54	2200	62	54	9.1 ± 0.8	20 ± 1	50 ± 3	3.9 ± 0.3	8.8 ± 0.8
16	18:47:47.02	-1:54:30.78	0.78 × 0.45	2200	68	0	25 ± 3	58 ± 4	21 ± 2	36 ± 6	90 ± 10
17	18:47:47.10	-1:54:27.07	0.51 × 0.45	1200	46	0	16 ± 6	18 ± 6	35 ± 2	6 ± 2	80 ± 30
18	18:47:46.25	-1:54:33.41	0.93 × 0.45	2600	59	0	22.7 ± 0.8	53 ± 1	23 ± 2	28 ± 4	39 ± 5
19	18:47:46.88	-1:54:25.74	0.45 × 0.45	1200	45	0	16 ± 5	19 ± 5	24 ± 2	9 ± 3	130 ± 40
20	18:47:45.29	-1:54:37.04	0.48 × 0.45	1200	47	0	13.2 ± 0.5	16.4 ± 0.6	23 ± 2	8 ± 1	110 ± 20
21	18:47:46.79	-1:54:16.06	0.47 × 0.45	1200	46	0	15 ± 1	20 ± 1	23 ± 2	10 ± 1	140 ± 20
22	18:47:47.05	-1:54:32.15	0.45 × 0.45	1200	40	0	13 ± 2	13 ± 2	21 ± 2	8 ± 2	100 ± 20
23	18:47:46.90	-1:54:24.30	0.63 × 0.45	1600	44	0	15 ± 5	20 ± 5	22 ± 2	11 ± 3	60 ± 20
24	18:47:46.97	-1:54:32.08	0.63 × 0.45	1600	29	0	10 ± 2	12 ± 2	21 ± 2	7 ± 1	37 ± 8
25	18:47:46.87	-1:54:14.62	0.45 × 0.45	1200	29	0	8 ± 1	8 ± 1	24 ± 2	4.0 ± 0.7	50 ± 10
26	18:47:44.94	-1:54:42.84	0.51 × 0.45	1200	28	0	7.1 ± 0.6	10.5 ± 0.7	23 ± 2	5.2 ± 0.7	70 ± 10
27	18:47:47.03	-1:54:24.91	0.62 × 0.45	1600	25	0	10 ± 5	11 ± 5	26 ± 2	5 ± 2	30 ± 10
28	18:47:44.40	-1:54:41.80	0.81 × 0.45	2300	22	0	5.8 ± 0.7	11.7 ± 0.8	23 ± 2	5.9 ± 0.9	12 ± 2
29	18:47:45.22	-1:54:38.81	0.45 × 0.45	1200	20	0	5.5 ± 0.5	6.6 ± 0.6	24 ± 2	3.1 ± 0.5	43 ± 6
30	18:47:47.18	-1:54:21.56	0.57 × 0.45	1400	19	26	5 ± 2	6 ± 2	62 ± 4	1.0 ± 0.3	10 ± 3
31	18:47:46.73	-1:54:17.53	0.59 × 0.45	1500	19	0	6.6 ± 0.9	8 ± 1	23 ± 2	4.3 ± 0.7	33 ± 6
32	18:47:46.57	-1:54:20.84	0.84 × 0.45	2400	20	0	7 ± 2	15 ± 2	28 ± 2	6 ± 1	11 ± 2
33	18:47:47.08	-1:54:28.80	0.72 × 0.46	2000	24	0	9 ± 4	17 ± 4	22 ± 2	9 ± 3	26 ± 8
34	18:47:46.52	-1:54:28.87	0.64 × 0.45	1700	18	0	6.9 ± 0.9	15 ± 1	25 ± 2	7 ± 1	35 ± 5
35	18:47:46.92	-1:54:31.75	0.73 × 0.46	2100	20	0	7 ± 3	15 ± 3	22 ± 2	8 ± 2	23 ± 5
36	18:47:46.64	-1:54:19.51	0.45 × 0.45	1200	18	0	5.8 ± 0.8	8 ± 1	27 ± 2	3.1 ± 0.5	43 ± 7
37	18:47:46.97	-1:54:12.96	0.73 × 0.45	2000	20	0	5 ± 1	11 ± 1	23 ± 2	5.2 ± 0.9	15 ± 3
38	18:47:47.02	-1:54:22.90	0.45 × 0.45	1200	16	0	5 ± 1	5 ± 2	24 ± 2	2.5 ± 0.8	30 ± 10
39	18:47:44.61	-1:54:42.16	0.47 × 0.45	1200	16	0	3.6 ± 0.6	4.8 ± 0.7	23 ± 2	2.3 ± 0.4	32 ± 6
40	18:47:46.35	-1:54:29.52	0.53 × 0.45	1200	17	0	5.8 ± 0.8	9.4 ± 0.9	24 ± 2	4.3 ± 0.7	59 ± 9
41	18:47:46.96	-1:54:23.87	0.68 × 0.45	1800	15	0	5 ± 3	7 ± 3	22 ± 2	4 ± 2	14 ± 7
42	18:47:44.78	-1:54:38.05	0.45 × 0.45	1200	13	0	3.0 ± 0.4	3.1 ± 0.4	24 ± 2	1.4 ± 0.2	19 ± 3
43	18:47:46.84	-1:54:31.64	0.64 × 0.46	1700	12	0	4 ± 3	5 ± 3	22 ± 2	2 ± 1	11 ± 7
44	18:47:44.76	-1:54:46.76	0.56 × 0.45	1300	15	0	3.2 ± 0.4	5.0 ± 0.4	22 ± 2	2.5 ± 0.4	28 ± 4
45	18:47:46.87	-1:54:36.50	0.45 × 0.45	1200	11	0	4.1 ± 0.9	4.2 ± 0.9	21 ± 2	2.4 ± 0.6	32 ± 8
46	18:47:46.77	-1:54:16.85	0.73 × 0.45	2000	12	0	4.7 ± 0.9	8.3 ± 0.9	21 ± 2	4.6 ± 0.8	14 ± 2
47	18:47:46.65	-1:54:31.90	0.81 × 0.59	2900	21	0	9 ± 2	35 ± 3	22 ± 2	19 ± 3	18 ± 3
48	18:47:46.82	-1:54:26.10	0.66 × 0.45	1700	12	0	5 ± 2	7 ± 2	23 ± 2	4 ± 1	16 ± 6
49	18:47:46.59	-1:54:20.48	0.73 × 0.48	2200	14	0	5 ± 1	9 ± 2	28 ± 2	3.7 ± 0.8	9 ± 2
50	18:47:46.97	-1:54:34.42	0.45 × 0.45	1200	11	0	3 ± 1	3 ± 1	21 ± 2	1.7 ± 0.8	20 ± 10

Supplementary Table 1 — Characteristics of the cores detected at 1.3 mm in W43-MM1 (continued).

#	RA [J2000]	Dec. [J2000]	Size ["×"]	FWHM [AU]	Sig	C_{line} [%]	$S_{1.3\text{mm}}^{\text{peak}}$ [mJy/beam]	$S_{1.3\text{mm}}^{\text{int}}$ [mJy]	\bar{T}_{dust} [K]	M_{core} [M_{\odot}]	\bar{n}_{H_2} $\times 10^7 \text{cm}^{-3}$
51	18:47:45.27	-1:54:40.03	0.96 × 0.52	3000	15	0	4.0 ± 0.7	10.2 ± 0.9	24 ± 2	4.8 ± 0.7	4.3 ± 0.6
52	18:47:46.80	-1:54:34.34	0.45 × 0.45	1200	10	0	3 ± 1	3 ± 1	21 ± 2	1.8 ± 0.8	20 ± 10
53	18:47:47.05	-1:54:18.90	0.77 × 0.55	2600	11	0	3.8 ± 0.7	7.5 ± 0.8	22 ± 2	4.0 ± 0.7	5.2 ± 0.9
54	18:47:45.06	-1:54:42.05	0.79 × 0.45	2200	11	0	2.9 ± 0.4	6.0 ± 0.5	23 ± 2	3.0 ± 0.4	7 ± 1
55	18:47:46.96	-1:54:35.42	0.63 × 0.45	1600	9.6	0	3 ± 1	3 ± 1	21 ± 2	2.0 ± 0.8	11 ± 4
56	18:47:47.00	-1:54:20.12	0.58 × 0.45	1400	11	0	3 ± 1	3 ± 1	21 ± 2	1.9 ± 0.9	17 ± 8
57	18:47:46.94	-1:54:19.37	0.45 × 0.45	1200	9.8	0	3 ± 1	3 ± 1	23 ± 2	1.7 ± 0.5	23 ± 8
58	18:47:47.08	-1:54:21.28	0.74 × 0.45	2000	11	0	4 ± 2	7 ± 3	25 ± 2	3 ± 1	8 ± 4
59	18:47:44.77	-1:54:44.17	0.63 × 0.45	1600	9.9	0	2.3 ± 0.9	3 ± 1	23 ± 2	1.4 ± 0.5	8 ± 3
60	18:47:44.76	-1:54:40.46	0.48 × 0.45	1200	10	0	2.4 ± 0.8	3.2 ± 0.9	23 ± 2	1.6 ± 0.5	22 ± 7
61	18:47:46.82	-1:54:35.53	0.7 × 0.53	2300	12	0	4 ± 1	10 ± 1	21 ± 2	6 ± 1	11 ± 2
62	18:47:45.79	-1:54:32.76	0.65 × 0.45	1700	11	0	3.7 ± 0.7	9 ± 1	24 ± 2	4.0 ± 0.6	19 ± 3
63	18:47:47.33	-1:54:12.82	0.47 × 0.45	1200	9.5	0	3.0 ± 0.7	3.8 ± 0.7	21 ± 2	2.1 ± 0.5	29 ± 7
64	18:47:46.99	-1:54:16.85	0.83 × 0.49	2600	10	0	3 ± 1	6 ± 1	22 ± 2	3.2 ± 0.7	5 ± 1
65	18:47:45.41	-1:54:37.44	0.56 × 0.45	1300	8.9	0	2.7 ± 0.6	3.1 ± 0.6	23 ± 2	1.6 ± 0.4	16 ± 4
66	18:47:47.00	-1:54:21.20	0.68 × 0.45	1800	9.0	0	3 ± 2	4 ± 2	22 ± 2	2.3 ± 0.9	9 ± 3
67	18:47:44.09	-1:54:48.89	0.66 × 0.47	1900	9.8	0	1.8 ± 0.3	4.7 ± 0.4	23 ± 2	2.2 ± 0.3	8 ± 1
68	18:47:47.02	-1:54:35.64	0.71 × 0.45	2000	8.7	0	3 ± 1	4 ± 1	21 ± 2	2.2 ± 0.6	7 ± 2
69	18:47:47.15	-1:54:20.05	0.5 × 0.45	1200	9.1	0	3 ± 1	5 ± 1	25 ± 2	2.1 ± 0.6	29 ± 8
70	18:47:47.20	-1:54:17.78	0.79 × 0.48	2400	9.1	0	3.3 ± 0.7	6.8 ± 0.8	23 ± 2	3.4 ± 0.6	6 ± 1
71	18:47:47.35	-1:54:13.46	0.62 × 0.45	1600	8.3	0	2.4 ± 0.6	3.1 ± 0.6	21 ± 2	1.7 ± 0.4	10 ± 2
72	18:47:46.65	-1:54:35.53	0.74 × 0.45	2100	8.2	0	3 ± 1	4 ± 1	20 ± 2	2.3 ± 0.8	6 ± 2
73	18:47:44.84	-1:54:42.95	0.63 × 0.53	2100	11	0	2.5 ± 0.8	6 ± 1	23 ± 2	2.8 ± 0.7	8 ± 2
74	18:47:45.17	-1:54:39.53	0.62 × 0.45	1600	8.0	0	2.2 ± 0.4	2.9 ± 0.5	24 ± 2	1.4 ± 0.3	8 ± 2
75	18:47:46.90	-1:54:35.10	0.73 × 0.45	2000	8.0	0	2 ± 1	3 ± 1	21 ± 2	1.9 ± 0.7	6 ± 2
76	18:47:46.93	-1:54:37.87	0.48 × 0.45	1200	8.0	0	2.8 ± 0.9	3 ± 1	21 ± 2	1.6 ± 0.6	22 ± 8
77	18:47:47.41	-1:54:25.60	0.73 × 0.45	2000	8.9	0	3.3 ± 0.9	6 ± 1	26 ± 2	2.6 ± 0.5	8 ± 2
78	18:47:46.43	-1:54:29.81	0.79 × 0.45	2200	7.7	0	2.3 ± 0.9	4 ± 1	23 ± 2	2.0 ± 0.6	5 ± 1
79	18:47:46.73	-1:54:25.63	0.8 × 0.46	2300	11	0	3 ± 1	7 ± 1	26 ± 2	2.8 ± 0.7	5 ± 1
80	18:47:47.15	-1:54:24.66	0.73 × 0.45	2000	7.2	0	3 ± 1	3 ± 1	22 ± 2	1.6 ± 0.7	5 ± 2
81	18:47:47.24	-1:54:18.47	0.74 × 0.46	2100	8.4	0	2.9 ± 0.6	5.4 ± 0.7	23 ± 2	2.6 ± 0.5	7 ± 1
82	18:47:46.90	-1:54:33.98	0.63 × 0.45	1600	7.7	0	3 ± 1	3 ± 1	21 ± 2	1.9 ± 0.6	10 ± 3
83	18:47:46.71	-1:54:24.19	0.75 × 0.45	2100	7.9	0	3 ± 1	4 ± 1	28 ± 2	1.5 ± 0.6	4 ± 2
84	18:47:46.37	-1:54:21.28	0.78 × 0.45	2200	9.7	0	3.1 ± 0.8	6 ± 1	26 ± 2	2.7 ± 0.5	6 ± 1
85	18:47:46.51	-1:54:26.42	0.7 × 0.5	2200	8.3	0	2.5 ± 0.9	5 ± 1	27 ± 2	1.8 ± 0.5	4 ± 1
86	18:47:46.74	-1:54:20.81	0.58 × 0.45	1400	7.4	0	2.5 ± 0.9	3.1 ± 0.9	26 ± 2	1.3 ± 0.4	11 ± 4
87	18:47:47.24	-1:54:16.06	0.54 × 0.45	1200	7.5	0	2.3 ± 0.6	2.7 ± 0.7	23 ± 2	1.4 ± 0.4	18 ± 5
88	18:47:47.28	-1:54:38.27	0.69 × 0.45	1900	7.4	0	2.5 ± 0.8	3.6 ± 0.8	21 ± 2	2.0 ± 0.5	7 ± 2
89	18:47:46.30	-1:54:28.76	0.68 × 0.45	1800	7.3	0	2.7 ± 0.9	3.6 ± 0.9	26 ± 2	1.5 ± 0.4	6 ± 2
90	18:47:46.39	-1:54:18.14	0.84 × 0.45	2400	7.2	0	2.7 ± 0.7	4.9 ± 0.8	24 ± 2	2.4 ± 0.5	4.4 ± 0.9
91	18:47:46.75	-1:54:33.30	0.63 × 0.45	1600	7.1	0	3 ± 1	3 ± 1	21 ± 2	1.8 ± 0.8	10 ± 4
92	18:47:46.86	-1:54:12.13	0.79 × 0.47	2300	7.9	0	2.5 ± 0.5	4.5 ± 0.5	22 ± 2	2.4 ± 0.4	4.7 ± 0.8
93	18:47:46.96	-1:54:40.86	0.69 × 0.45	1900	7.0	0	2.7 ± 0.7	3.7 ± 0.8	21 ± 2	2.1 ± 0.5	8 ± 2
94	18:47:46.74	-1:54:37.80	0.67 × 0.45	1800	7.0	0	2.7 ± 0.6	3.9 ± 0.7	21 ± 2	2.2 ± 0.5	9 ± 2
95	18:47:46.58	-1:54:25.67	0.72 × 0.45	2000	7.1	0	2.7 ± 0.9	3.1 ± 0.9	28 ± 2	1.2 ± 0.4	4 ± 1
96	18:47:47.08	-1:54:23.54	0.65 × 0.45	1700	7.4	0	3 ± 1	4 ± 1	24 ± 2	2.0 ± 0.6	9 ± 3
97	18:47:47.16	-1:54:10.15	0.72 × 0.45	2000	7.6	0	1.9 ± 0.4	4.1 ± 0.6	21 ± 2	2.3 ± 0.5	7 ± 1
98	18:47:46.40	-1:54:28.37	0.68 × 0.52	2200	7.0	0	2.3 ± 0.9	4 ± 1	27 ± 2	1.5 ± 0.4	3 ± 1
99	18:47:45.39	-1:54:39.13	1.26 × 0.55	3900	8.5	0	1.7 ± 0.6	7.1 ± 0.9	23 ± 2	3.4 ± 0.6	1.4 ± 0.2
100	18:47:45.22	-1:54:40.46	0.7 × 0.47	2100	6.8	0	2.1 ± 0.5	3.1 ± 0.5	24 ± 2	1.4 ± 0.3	4.1 ± 0.8

Supplementary Table 1 — Characteristics of the cores detected at 1.3 mm in W43-MM1 (continued).

#	RA [J2000]	Dec. [J2000]	Size ["×"]	FWHM [AU]	Sig	C_{line} [%]	$S_{1.3\text{mm}}^{\text{peak}}$ [mJy/beam]	$S_{1.3\text{mm}}^{\text{int}}$ [mJy]	\bar{T}_{dust} [K]	M_{core} [M_{\odot}]	\bar{n}_{H_2} $\times 10^7 \text{cm}^{-3}$
101	18:47:47.30	-1:54:35.03	0.59 × 0.45	1500	6.8	0	2.8 ± 0.7	4.1 ± 0.8	21 ± 2	2.3 ± 0.6	17 ± 4
102	18:47:47.15	-1:54:33.19	0.7 × 0.45	1900	6.7	0	2 ± 1	2 ± 1	21 ± 2	1.2 ± 0.6	4 ± 2
103	18:47:46.74	-1:54:19.44	0.51 × 0.45	1200	7.0	0	2 ± 1	2 ± 1	26 ± 2	1.1 ± 0.5	15 ± 6
104	18:47:46.80	-1:54:18.43	0.61 × 0.45	1500	6.9	0	2.2 ± 0.8	2.9 ± 0.9	24 ± 2	1.3 ± 0.4	9 ± 3
105	18:47:46.50	-1:54:16.24	0.46 × 0.45	1200	6.9	0	1.9 ± 0.6	1.9 ± 0.6	23 ± 2	1.0 ± 0.3	13 ± 4
106	18:47:46.40	-1:54:36.61	0.63 × 0.45	1600	6.9	0	2.3 ± 0.6	3.6 ± 0.7	21 ± 2	2.0 ± 0.5	11 ± 3
107	18:47:47.24	-1:54:35.60	0.57 × 0.47	1500	6.8	0	2.6 ± 0.7	3.9 ± 0.7	21 ± 2	2.2 ± 0.5	15 ± 3
108	18:47:45.32	-1:54:39.13	0.69 × 0.45	1900	6.6	0	1.9 ± 0.9	2.6 ± 0.9	24 ± 2	1.2 ± 0.5	4 ± 2
109	18:47:46.60	-1:54:18.00	0.67 × 0.45	1800	6.7	0	2.3 ± 0.9	4 ± 1	25 ± 2	1.7 ± 0.5	7 ± 2
110	18:47:47.16	-1:54:11.30	0.65 × 0.45	1700	6.8	0	2.1 ± 0.6	3.4 ± 0.6	21 ± 2	1.9 ± 0.4	9 ± 2
111	18:47:46.80	-1:54:20.16	0.7 × 0.45	1900	6.5	0	2 ± 1	3 ± 1	25 ± 2	1.4 ± 0.6	5 ± 2
112	18:47:46.30	-1:54:37.37	0.66 × 0.45	1800	6.4	0	2.3 ± 0.5	2.7 ± 0.5	22 ± 2	1.5 ± 0.3	7 ± 2
113	18:47:47.03	-1:54:10.58	0.64 × 0.47	1800	6.4	0	1.5 ± 0.5	2.2 ± 0.6	21 ± 2	1.2 ± 0.4	5 ± 2
114	18:47:46.91	-1:54:17.75	0.53 × 0.45	1200	6.4	0	1.9 ± 0.5	2.2 ± 0.5	22 ± 2	1.1 ± 0.3	16 ± 4
115	18:47:46.76	-1:54:12.64	0.77 × 0.57	2700	6.9	0	2.4 ± 0.2	5.3 ± 0.2	22 ± 2	2.8 ± 0.4	3.5 ± 0.5
116	18:47:46.18	-1:54:30.92	0.62 × 0.45	1600	6.4	0	2.2 ± 0.6	3.5 ± 0.7	23 ± 2	1.7 ± 0.4	10 ± 2
117	18:47:46.67	-1:54:23.80	0.97 × 0.47	2800	6.7	0	3 ± 1	6 ± 1	29 ± 2	2.3 ± 0.6	2.6 ± 0.6
118	18:47:47.04	-1:54:14.90	0.72 × 0.45	2000	6.1	0	1.6 ± 0.7	2.5 ± 0.7	23 ± 2	1.2 ± 0.4	4 ± 1
119	18:47:47.37	-1:54:11.81	0.54 × 0.45	1200	6.4	0	1.8 ± 0.5	2.4 ± 0.6	21 ± 2	1.4 ± 0.4	18 ± 5
120	18:47:46.80	-1:54:11.92	0.76 × 0.63	2900	6.2	0	2.5 ± 0.4	5.7 ± 0.4	22 ± 2	3.0 ± 0.5	2.9 ± 0.5
121	18:47:47.00	-1:54:39.53	0.74 × 0.45	2100	6.3	0	1.7 ± 0.8	2.4 ± 0.9	21 ± 2	1.3 ± 0.5	4 ± 1
122	18:47:44.76	-1:54:53.14	0.59 × 0.56	2000	6.7	0	1.3 ± 0.4	3.1 ± 0.6	23 ± 2	1.5 ± 0.3	4.4 ± 0.9
123	18:47:46.99	-1:54:4.90	0.85 × 0.45	2400	6.3	0	1.7 ± 0.4	3.2 ± 0.5	21 ± 2	1.8 ± 0.4	3.3 ± 0.7
124	18:47:47.37	-1:54:21.38	0.84 × 0.45	2300	5.8	0	2.4 ± 0.7	5.0 ± 0.8	25 ± 2	2.2 ± 0.4	4.1 ± 0.8
125	18:47:47.47	-1:54:38.09	0.63 × 0.45	1700	6.3	0	2.2 ± 0.5	3.3 ± 0.6	23 ± 2	1.6 ± 0.4	9 ± 2
126	18:47:44.57	-1:54:39.71	0.83 × 0.53	2700	6.4	0	1.5 ± 0.5	4.3 ± 0.8	23 ± 2	2.1 ± 0.5	2.6 ± 0.6
127	18:47:46.25	-1:54:22.25	1.02 × 0.55	3300	6.2	0	1.9 ± 0.6	5.8 ± 0.9	27 ± 2	2.4 ± 0.4	1.6 ± 0.3
128	18:47:47.29	-1:54:26.32	0.88 × 0.57	3000	5.9	0	1.7 ± 0.9	4 ± 1	23 ± 2	2.0 ± 0.6	1.7 ± 0.5
129	18:47:46.56	-1:54:6.95	0.87 × 0.48	2600	6.1	0	1.5 ± 0.4	3.7 ± 0.5	23 ± 2	1.8 ± 0.3	2.6 ± 0.5
130	18:47:46.37	-1:54:47.74	0.98 × 0.45	2700	5.7	0	1.9 ± 0.4	4.5 ± 0.5	21 ± 2	2.5 ± 0.5	3.0 ± 0.5
131	18:47:46.55	-1:54:16.63	1.11 × 0.5	3300	5.8	0	1.6 ± 0.6	5.1 ± 0.8	23 ± 2	2.6 ± 0.5	1.7 ± 0.3

Notes: RA, right ascension; Dec., declination; Size at half-maximum; FWHM, Size deconvolved with the 0.44'' telescope beam; Sig, significance level of *getsources* core extraction; line contamination ratio, $C_{\text{line}} = [S^{\text{peak}}(1.9 \text{ GHz}) - S^{\text{peak}}(65 \text{ MHz})] / S^{\text{peak}}(65 \text{ MHz})$; $S_{1.3\text{mm}}^{\text{peak}}$ and $S_{1.3\text{mm}}^{\text{int}}$, peak and total integrated flux corrected for line contamination and local background; \bar{T}_{dust} , mean dust temperature read in Fig. 3; M_{core} , core mass; mean density, $\bar{n}_{\text{H}_2} = \frac{M_{\text{core}}}{\frac{4}{3}\pi\mu m_{\text{H}}(FWHM/2)^3}$, where $\mu = 2.8$ is the mean molecular weight per H_2 molecule and m_{H} is the hydrogen mass.

Supplementary Table 2 — Tests performed to evaluate the uncertainty of the reference CMF fit of Fig. 2b.

	Mass range	γ
Reference cumulative CMF of all cores extracted by <i>getsources</i> with 5σ uncertainty derived from the mass uncertainties low-mass regime (high-mass regime, 9 cores) with a lower completeness level	$> 1.6 M_{\odot}$	-0.96 ± 0.02
	$> 1.6 M_{\odot}$	-0.96 ± 0.13
	$1.6 - 20 M_{\odot}$	-0.93 ± 0.02
	$(> 20 M_{\odot})$	(-1.3 ± 0.3)
	$> 4.5 M_{\odot}$	-0.99 ± 0.04
CMF of the 94 most robust cores	$> 1.6 M_{\odot}$	-0.90 ± 0.02
CMF with core masses estimated in the optical thin approximation	$> 1.6 M_{\odot}$	-0.98 ± 0.04
Differential CMF with all cores and default assumptions	$> 1.6 M_{\odot}$	-0.90 ± 0.06
CMF built from cores extracted in a classic-cleaned image in a merged (7 m + 12 m) image with MRE-GAUSSCLUMPS	$> 1.6 M_{\odot}$	-1.10 ± 0.05
	$> 1.6 M_{\odot}$ or $> 5 M_{\odot}$	-1.10 ± 0.04
	$> 1.6 M_{\odot}$ or $> 5 M_{\odot}$	-1.08 ± 0.04

Notes: CMFs are fitted by power-laws of the form $N(> \log(M)) \propto M^{\gamma}$, except for the differential CMF where the power-law is $dN/d \log(M) \propto M^{\gamma}$. Several mass ranges are used to fit the CMFs of less-constrained core samples derived from the merged (7 m + 12 m) image and the MRE-GaussClumps algorithm. Except when specified otherwise, all uncertainties given here are 1σ .